

## Neutron Band in Solids

Hideo KOZIMA

*Department of Physics, Faculty of Science, Shizuoka University,  
 836 Oya, Shizuoka 422-5829*

(Received May 13, 1998)

KEYWORDS: metals, surfaces and interfaces, neutron energy spectrum, neutron band

Low energy neutron with a de Broglie wave length comparable with lattice constants is widely used as a tool for structural analysis of matter.<sup>1,2)</sup> Characteristic behaviors of neutron have been investigated in these several tens years.

The neutrons interacting with nuclei bound in crystals (lattice nuclei)<sup>3,4)</sup> have exhibited such a similar feature of interaction with individual nucleus as in vacuum (free nucleus).

On the other hand, Bragg reflection due to the interaction of a neutron wave with a lattice of nuclei has been used to trap neutrons between two single crystal plates of Si.<sup>5)</sup> Also, a shift of effective mass of neutrons with energies near the Bragg condition in a crystal was shown<sup>6)</sup> by an experiment of neutron propagation through the crystal under diffraction condition. Recently, artificial periodic structures<sup>7,8)</sup> of a potential on the cold neutron have been used in neutron physics to investigate the quantum precession and the splitting of neutron spin, tunneling time of a potential barrier and trapping time in potential well in addition to the traditional diffraction and reflection just similar to those for the electron.

Furthermore, a change of decay behavior of neutrons in strong electromagnetic field was predicted by a calculation.<sup>9)</sup> There are a new field of research in the decay behavior of neutron interacting with other nuclei through electromagnetic and strong interactions.

This paper gives a simple treatment of band structure of low energy (thermal) neutrons in a crystal (neutron band) and a possible trapping mechanism of neutrons related with reflection at a boundary due to a difference of band structures in neighboring crystals. A numerical result is shown by a simple one-dimensional Kronig-Penny model with Fermi quasi-potential, i.e. a delta-function type potential. Possible physics of the system composed of a crystal and the trapped thermal neutron is discussed.

A thermal neutron with de Broglie wave length comparable with the lattice constant of a crystal behaves as a wave and interacts with nuclei in the lattice (lattice nuclei) through the nuclear force. The interaction could be treated approximately by Fermi pseudo-potential (zero-radius potential<sup>10)</sup>)

$$V(r) = \frac{2\pi\hbar^2 b}{m} \frac{\delta(r)}{r^2} \quad (1)$$

with  $b$  a parameter with dimension of length (an effective scattering length). The wave function of the neutron with mass  $m$  will be calculated with Wigner-Seitz

method<sup>11)</sup> solving an equation

$$\left\{ -\frac{\hbar^2}{2mr^2} \frac{d}{dr} \left( r^2 \frac{d}{dr} \right) + V(r) \right\} \psi(r) = E\psi(r) \quad (2)$$

subject to a boundary condition

$$\left( \frac{\partial \psi}{\partial r} \right) = 0, \quad (\text{at } r = r_0) \quad (3)$$

where  $r_0$  is the radius of the so-called s-sphere having the same volume as the atomic volume of the crystal.

To illustrate the essential point of this treatment, we calculate the energy band structure of a particle with mass  $m$  in the one dimensional Kronig-Penny lattice with a lattice constant  $a$  and an attractive delta-function potential:

$$V(x) = -\frac{\hbar^2 P}{ma} \delta(x + na), \quad (n = 0, \pm 1, \pm 2, \dots) \quad (4)$$

where  $P$  is a constant proportional to the intensity of the attractive potential (width  $\times$  depth). The energy  $E$  of the neutron with wave vector  $k$  in this crystal is determined by an equation<sup>11)</sup>

$$-P \frac{\sin \alpha a}{\alpha a} + \cos \alpha a = \cos ka \quad (5)$$

where  $\alpha = (2mE/\hbar^2)^{1/2}$ . Band structure of this energy spectrum was calculated numerically and plotted in Fig. 1 in the reduced zone scheme ( $-\pi \leq ka \leq \pi$ ). The value of  $P$  was assumed arbitrarily as  $3\pi/2$  which corresponds to a square well potential with a depth  $2 \times 10^3$  eV and a width  $10^{-13}$  cm considering the interaction through the nuclear force. Increase of the parameter  $P$  makes the width of the allowed band narrower. The energy of the edge of the lowest energy band is inversely proportional to the square of the lattice constant  $a^2$  and is  $2.04 \times 10^{-2}$  eV for  $a = 10^{-8}$  cm. It is interesting to notice that the energy of this band edge is somewhat lower than the thermal energy at room temperature (300 K) for a lattice with a lattice constant  $2 \sim 3$  Å.

On the other hand, the wave function of the neutron in the band has a large amplitude at lattice points where are centers of the attractive potential. So, the neutron in a crystal has a band structure in its energy spectrum and localized probability amplitude at nuclei on the lattice points.

The neutron band illustrated in Fig. 1 will contribute to trapping of thermal neutrons in a crystal if the crystal is surrounded by a wall with a different band structure. A neutron in the crystal can not stay in the wall if the energy of the neutron corresponds to a forbidden energy of the band in the wall. Then in such a situation, a neutron with a thermal or epi-thermal energy can exist trapped in the lowest neutron band in the crystal if the width of the wall is thick enough to prevent tunneling.

The neutron band in a three-dimensional lattice could be calculated solving eq. (2) with the potential eq. (1) and the boundary condition eq. (3) and is a future program though the essential feature of the neutron band is illustrated above for the one dimensional case.

As the position and width of the allowed band depend on the lattice constant  $a$  and the parameter  $P$  (proportional




Fig. 1. Energy band structure of neutron in one dimensional attractive delta-function type Kronig-Penny potential. Parameter  $P$  is taken arbitrarily as  $3\pi/2$  and the wave number is reduced to  $|ka| \leq \pi$ . For  $a = 10^{-8}$  cm, the energy of the edge of the lowest band is  $2 \times 10^{-2}$  eV.

tional to strength of potential), a neutron in a band in one region could be reflected at the boundary of the adjacent region with different lattice constant and/or constituent nuclei. This situation occurs, for instance, in palladium hydrides (deuterides) with periodic distribution of occluded hydrogen (deuteron) or with a surface layer of alkali (Li) metal on the surface.

The Kronig-Penny model of a potential on the cold neutron has been formed artificially with a laminar structure of metals<sup>7,8</sup>) and the artificial periodic potential thus formed has been used to diffract and reflect neutrons for investigation in neutron physics. In this case, however, the potential is not that given in eq. (1) but one averaged over nuclei in a layer of the structure to form a diffractive index for the neutron.

The notion of a neutron band in solids proposed in this paper for the first time is a novel one corresponding to the popular electron band in solids and the photon band in artificial periodic structures. It is interesting to notice that these all particles are used in structural analysis as electron, X-ray and neutron diffractions widely and complementally in science and technology now.

On the other hand, those band structures in the energy spectra have fairly different characters. The electron band, as we know well, had been worked out in 1930's and explained successfully characteristics of solids and had been the basis of the modern technology. The photon band has recently realized using artificial materials with an appropriate period for the wave length of a relevant photon.

Recently, multi-layer structures of crystals, for instance Permalloy(100 Å)/ Ge(800 Å)/ Permalloy(100 Å), have been used to investigate trapping and tunneling of a cold neutron.<sup>8)</sup> This is a new direction of low energy

neutron physics where the property of a neutron in a potential is an object of research instead of the external effect of a crystal or a potential structure on the neutron investigated hitherto.<sup>1-7)</sup>

There are several possibilities to find out characteristics of the energy band of thermal neutrons due to the nuclear interaction not found in electron and photon bands due to electrostatic and electromagnetic interactions, respectively. The characteristics of the cold neutrons in the multi-layer films are rather similar to the electron and photon band and those of the band structure investigated in this paper will be different from them.

One of characteristics of the neutron band, if we look into details, calculated by the Kronig-Penny model and given in Fig. 1 is the shape of the lowest band. The energy minimum of this neutron band is at  $k = \pi/a$  in contrast to at  $k = 0$  in the case of the electron band. This characteristic shape of the lowest band of neutron energy spectrum results in existence of trapped neutrons with opposite momenta and spins even when the concentration of the trapped neutron is very low.

If two trapped neutrons with opposite lattice momenta and spins form a Cooper pair (neutron Cooper pair) by the interaction through phonons, the energy of the system becomes lower to make the life time of the neutron elongated further.

Behaviors of thermal neutrons in quiet solids have not attracted attention except those works on the external effects of crystals for the neutron cited above<sup>3-8)</sup> though there are interesting features of solid state-nuclear physics pointed out above as 1) elongation of neutron life time, 2) destabilization of foreign nuclei in a lattice, and 3) neutron fusion with or scattering by a nucleus exerting large perturbation on it, which may be more or less interesting than those in high energy neutron physics developed in relation with nuclear reactors.

- 1) C. G. Shull and E. O. Wollan: *Solid State Physics*, ed. F. Seitz and D. Turnbull (Academic Press, New York, 1956) Vol. 2, p. 1337.
- 2) W. G. Williams: *Polarized Neutron, Oxford Series on Neutron Scattering in Condensed Matter, No. 1* (Oxford University Press, Oxford, England, 1957).
- 3) A. Michaudon, H. Derrien, C. Le Pipec and P. Ribon: *Comptes Rendus* **255** (1962) 2086.
- 4) H. E. Jackson and J. E. Lynn: *Phys. Rev.* **127** (1962) 461.
- 5) M. Shuster, H. Rauch, E. Seidl, E. Jericha and C. J. Carlile: *Phys. Lett. A* **144** (1990) 297.
- 6) A. Zeilinger, C. G. Shull, M. A. Horne and K. D. Finkelstein: *Phys. Rev. Lett.* **57** (1986) 3089.
- 7) T. Ebisawa, H. Hunabashi, S. Tazaki, Y. Otake, T. Kawai, M. Hino, N. Achiwa and T. Akiyoshi: *J. Neutron Research* **4** (1995) 157.
- 8) N. Achiwa, M. Hino, S. Tazaki, T. Ebisawa, T. Akiyoshi and T. Kawai: *Physica B* (Proc. ICNS97, Toronto, 1998) to be published.
- 9) I. M. Ternov, V. N. Dodionov, V. G. Zhulego and A. I. Studenikin: *Sov. J. Nucl. Phys.* **28** (1979) 747.
- 10) Y. N. Demkov and V. N. Ostrovskii: *Methods of Zero-radius Potential in Atomic Physics* (Leningrad Univ., Press, Leningrad, 1975) [in Russian].
- 11) C. Kittel: *Introduction to Solid State Physics* (John Wiley, New York, 1986).